arXiv:0910.0743v1 [math.DS] 5 Oct 2009

DYNAMICAL APPROXIMATION AND KERNELS OF NONESCAPING-HYPERBOLIC COMPONENTS

HELENA MIHALJEVIĆ-BRANDT

Abstract. Let $\mathscr{F}_n$ be families of entire functions, holomorphically parametrized by a complex manifold M. We consider those parameters in M that correspond to *nonescaping-hyperbolic* functions, i.e., those maps $f \in \mathscr{F}_n$ for which the postsingular set $P(f)$ is a compact subset of the Fatou set $\mathcal{F}(f)$ of f. We prove that if $\mathscr{F}_n \to \mathscr{F}_\infty$ in the sense of a certain dynamically sensible metric, then every nonescaping-hyperbolic component in the parameter space of $\mathscr{F}_\infty$ is a *kernel* of a sequence of nonescaping-hyperbolic components in the parameter spaces of $\mathscr{F}_n$. Parameters belonging to such a kernel do not always correspond to hyperbolic functions in $\mathscr{F}_\infty$. Nevertheless, we show that these functions must be J-stable. Using quasiconformal equivalences, we are able to construct many examples of families to which our results can be applied.

1. Introduction

Let $\mathrm{Hol}^*(\mathbb{C})$ be the space of all entire functions that are not constant or linear. A map $f \in \mathrm{Hol}^*(\mathbb{C})$ is said to be *nonescaping-hyperbolic* if its postsingular set $P(f)$ is a compact subset of the Fatou set $\mathcal{F}(f)$. (Basic definitions and notations are revised in Section 2.) We want to understand how nonescaping-hyperbolic maps behave under certain small perturbations. It is known that within the space of all polynomials or the space of all transcendental entire maps with finitely many singular values, (nonescaping-)hyperbolic functions exhibit particularly simple and stable dynamics [17, 7].

The space $\mathrm{Hol}^*(\mathbb{C})$ is naturally equipped with the topology of locally uniform convergence. However, this topology is not convenient for our dynamical considerations since maps that are nearby in the corresponding metric often have completely different dynamics (see e.g. Example 2). We introduce a new metric χ_{dyn} on $\mathrm{Hol}^*(\mathbb{C})$ which is dynamically more sensible: two entire maps are close with respect to χ_{dyn} if their locally uniform distance *and* the Hausdorff distance between their sets of singular values is small.

There are certainly many examples of entire maps that converge in the metric χ_{dyn} to a map in $\mathrm{Hol}^*(\mathbb{C})$; the probably most prominent is the approximation of the exponential map $E(z) = \mathrm{e}^z$ by the polynomials $P_n(z) = (1 + z/n)^n$. In the article [4], Devaney, Goldberg and Hubbard investigated the relationship between the families $E_\lambda(z) = \lambda\,\mathrm{e}^z$ and $P_{n,\lambda}(z) = \lambda\,(1 + z/n)^n$. The result was a nice connection between the respective parameter spaces: the authors proved pointwise convergence of nonescaping-hyperbolic components (i.e. connected components of the set of those $\lambda \in \mathbb{C}$ for which E_λ or $P_{d,\lambda}$, respectively, is nonescaping-hyperbolic) as well as convergence of certain external rays to curves called "hairs" in exponential parameter space (see also [5]). One can say that this point of view has provided an important conceptional basis for much subsequent work on the exponential family.

In this article, we embed the underlying approximation idea into a general setup. More precisely, let M be a complex manifold and let $\chi_{M'}$ be a metric on $M' := (\mathbb{N} \cup \{\infty\}) \times M$. Let $\mathrm{Hol}^*_b(\mathbb{C})$ denote the set of all maps in $\mathrm{Hol}^*(\mathbb{C})$ with bounded sets of singular values. Our main result is the following.

Theorem 1.1. *For every $n \in \mathbb{N} \cup \{\infty\}$ let $\mathscr{F}_n = \{f_{n,\lambda}\} \subset \mathrm{Hol}^*_b(\mathbb{C})$ be a family of entire functions that depend holomorphically on $\lambda \in M$. Assume that for every n, the singular values of all maps in $\mathscr{F}_n$ are holomorphically parametrized by M, and that $F : M' \to \mathrm{Hol}^*_b(\mathbb{C}), (n, \lambda) \mapsto f_{n,\lambda}$ is continuous with respect to the metrics $\chi_{M'}$ and χ_{dyn}.*

If $\widetilde{H}$ is a kernel of a sequence of hyperbolic components of $\mathscr{F}_n$, then exactly one of the following statements holds:

- *(i) The map $f_\lambda \in \mathscr{F}_\infty$ is not hyperbolic for any $\lambda \in \widetilde{H}$.*
- *(ii) There is a hyperbolic component H_∞ of $\mathscr{F}_\infty$ such that $\widetilde{H} = H_\infty$.*

Remark. Our result is a natural generalization of a theorem by Krauskopf and Kriete, who considered holomorphic families $\mathscr{F}_n = \{f_\lambda : \lambda \in \mathbb{C}\}$, $n \in \mathbb{N} \cup \{\infty\}$, of entire maps for which the sets of singular values are holomorphically parametrized and for which there exists an integer N with $|S(f_{n,\lambda})| = N < \infty$ for all $n \in \mathbb{N} \cup \{\infty\}$ and all $\lambda \in \mathbb{C}$. They proved the same conclusions as in Theorem 1.1, provided that $\mathscr{F}_n \to \mathscr{F}_\infty$ uniformly on compact subsets of $\mathbb{C} \times \mathbb{C}$ [13]. Our proof of the above Theorem follows the same idea as in [13].

The first case in Theorem 1.1 does indeed occur; an example is given in Section 3. Nevertheless, parameters that belong to a kernel define maps which exhibit certain stability. Under the same assumptions (and notations) as in Theorem 1.1 we prove the following result.

Theorem 1.2. *Let λ belong to a kernel $\widetilde{H}$. Then $f_\lambda \in \mathscr{F}_\infty$ is a J-stable function.*

We define J-stability analogously to the classical definition for rational maps (see Definition 3.4).

Given an entire function f together with a sequence of entire maps f_n for which $\chi_{\mathrm{dyn}}(f_n, f) \to 0$ when $n \to \infty$, there is a natural way – using quasiconformal equivalence classes – to construct suitable holomorphic families $\mathscr{F}_n \ni f_n$ that satisfy the assumptions of Theorem 1.1. For a lot of explicit functions, including many examples that have been of particular dynamical interest in the last decades, non-trivial approximations in the sense of χ_{dyn} are known. However, for an *arbitrary* (transcendental) entire map there seems to be no general concept of how to establish a non-trivial sequence of maps f_n such that $\chi_{\mathrm{dyn}}(f_n, f) \to 0$. This leaves open an interesting function-theoretic question.

Structure of the article. In Section 2 we introduce basic definitions, notations and preliminary concepts. Section 3 addresses the proofs of Theorems 1.1 and 1.2. In the final part, we discuss examples to which our results can be applied.

1.1. **Acknowledgements.** I want to thank Lasse Rempe for his encouragement to extend the results from my Diploma thesis and for his continual help and support. I would also like to thank Hartje Kriete for introducing me to the subject during my time as a graduate student in Göttingen.

2. PRELIMINARY CONSTRUCTIONS

We denote the complex plane by $\mathbb{C}$ and the Riemann sphere by $\widehat{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$. For a subset V of a metric space M we denote the ε-neighbourhood of V by $U_\varepsilon(V)$. We will write $A \Subset B$ if A is a relatively compact subset of B, i.e., if $\overline{A}$ is compact and contained in B. If not stated differently, the boundary ∂A and the closure $\overline{A}$ of a set $A \subset \mathbb{C}$ is always understood to be taken relative to the complex plane. Throughout this article, we will mainly consider nonconstant, nonlinear entire functions $f : \mathbb{C} \to \mathbb{C}$, hence f will either be a polynomial of degree ≥ 2 or a transcendental entire map. Recall that the space of all such maps is denoted by $\mathrm{Hol}^*(\mathbb{C})$.

We denote by $C(f) := \{w \in \mathbb{C} : \exists z : f^{'}(z) = 0 \text{ and } f(z) = w\}$ the set of all critical values and by $A(f)$ the set of all (finite) asymptotic values of f. Recall that a point $w \in \mathbb{C}$ is an asymptotic value of f if there is a path to ∞ along which f converges to w. Note that a polynomial has no asymptotic values. The set $S(f)$ of *singular values* of f is defined to be the smallest closed set in $\mathbb{C}$ such that $f : \mathbb{C} \setminus f^{-1}(S(f)) \to \mathbb{C} \setminus S(f)$ is a covering map. It is well-known that $S(f)$ can be written as $S(f) = \overline{C(f) \cup A(f)}$ (see e.g. [9, Lemma 1.1]). We define

$$\mathrm{Hol}_b^*(\mathbb{C}) := \{f \in \mathrm{Hol}^*(\mathbb{C}) : S(f) \text{ is bounded}\}$$

to be the set of all nonconstant, nonlinear entire maps with bounded singular sets. Finally, we denote the *postsingular set* of f by $P(f) := \overline{\bigcup_{n \geq 0} f^n(S(f))}$.

Let $w \in \mathbb{C}$ be a periodic point of f of period k, that is, $f^k(w) = w$ and $k > 0$ is minimal with this property. The set $O(w) := \{w, f(w), ..., f^{k-1}(w)\}$ is called the (forward) *orbit* or *cycle* of w. The *multiplier* $\mu(w)$ of w is defined as $\mu(w) := (f^k)^{'}(w)$. We call w *attracting* if $|\mu(w)| < 1$, *repelling* if $|\mu(w)| > 1$ and *indifferent* if $|\mu(w)| = 1$. An attracting periodic point w with $\mu(w) = 0$ is called *superattracting*. If w is a periodic point then $\mu(f^i(w)) = \mu(w)$ for all i, hence we can extend the classification of w to the whole cycle $O(w)$ and speak of the multiplier of the cycle etc.

If O is an attracting periodic cycle of f of period n, then the *basin of attraction* or *attracting basin of* O is the open set $A(O)$ consisting of all points z for which the successive iterates $f^n(z), f^{2n}(z), \ldots$ converge to some point of O. If $w \in O$ then the component $A^*(w)$ of $A(O)$ that contains w is called the *immediate basin of attraction of* w. (The immediate basin of the cycle O is then the union of the immediate basins of the points in O.) We denote the set of all points that converge to an attracting cycle of f by $\mathcal{A}(f)$.

Let $f \in \mathrm{Hol}^*(\mathbb{C})$. The objects of main dynamical interest related to f are the *Fatou set* $\mathcal{F}(f)$ of f, defined as the set of all points that have a neighbourhood in which the iterates $(f^n)_{n \in \mathbb{N}}$ form a normal family in the sense of Montel, and its complement the *Julia set* $\mathcal{J}(f) := \mathbb{C} \setminus \mathcal{F}(f)$ of f.

Remark. A polynomial f extends naturally to a map $\widehat{f}:\widehat{\mathbb{C}}\to\widehat{\mathbb{C}}$ with $\widehat{f}(\infty)=\infty$. In this case, ∞ is a critical point and a critical value of $\widehat{f}$. However, the dynamically relevant sets are deliberately chosen to be subsets of the plane (rather than the sphere) since the consideration of the point at ∞ does not contribute to the understanding of the dynamics: for every transcendental entire map, ∞ is an essential singularity, while for a polynomial, ∞ is its only preimage.

The *escaping set* of f is defined to be

$$I(f):=\{z\in\mathbb{C}:\lim_{n\to\infty}f^n(z)=\infty\}.$$

A point $z\in I(f)$ is called an *escaping point*. For a polynomial f (of degree ≥ 2), the escaping set belongs to $\mathcal{F}(f)$, since the point at ∞ can be considered as a superattracting fixed point of the holomorphic extension of f to $\widehat{\mathbb{C}}$ and the escaping set as its basin of attraction. In contrast, if $f\in\operatorname{Hol}^*_b(\mathbb{C})$ is transcendental entire, then the relation $\mathcal{J}(f)=\overline{I(f)}$ holds [6, p. 344].

For more background in holomorphic dynamics, we refer to the expository works [1, 18]. The classification of components of the Fatou set of an entire map will be used implicitly throughout this article and can be found in [1].

Nonescaping-hyperbolic maps. Let us start with the definitions of hyperbolic and nonescaping-hyperbolic functions.

Definition 2.1. A map $f\in\operatorname{Hol}^*(\mathbb{C})$ is called *hyperbolic* if $S(f)$ is bounded and $P(f)\subset\mathcal{F}(f)$. We say that f is *nonescaping-hyperbolic* if $P(f)$ is a bounded subset of $\mathcal{F}(f)$.

Clearly, every nonescaping-hyperbolic map is also hyperbolic. Recall that the set $S(f)$ is closed by definition, hence if f is hyperbolic then $S(f)$ is a compact subset of $\mathcal{F}(f)$. Similarly for $P(f)$ when f is nonescaping-hyperbolic.

The next observation will be used frequently; it follows mainly from classical results in holomorphic dynamics but we could not localize a reference, hence we include a proof for completeness.

Proposition 2.2. *Let $f\in\operatorname{Hol}^*(\mathbb{C})$. Then f is nonescaping-hyperbolic if and only if $S(f)$ is a bounded subset of $\mathcal{A}(f)$.*

Let f be nonescaping-hyperbolic. Then $\operatorname{Attr}(f)$ is finite, and if f is transcendental entire then $\mathcal{F}(f)=\mathcal{A}(f)\neq\emptyset$, otherwise $\mathcal{F}(f)=\mathcal{A}(f)\cup I(f)$.

Proof. The statement is well-known for polynomials [18, Theorem 19.1] hence we can restrict to transcendental entire maps.

Assume that $S(f)$ is bounded and every $w\in S(f)$ converges to an attracting cycle of f. Then the components of $\mathcal{A}(f)$ form an open cover of $S(f)$ and since $S(f)$ is compact, there is a finite subcover. In particular, it follows that $P(f)$ is a compact subset of $\mathcal{A}(f)\subset\mathcal{F}(f)$, hence f is nonescaping-hyperbolic.

Now assume that $P(f)$ is a compact subset of $\mathcal{F}(f)$ and let U be a component of $\mathcal{F}(f)$. Then the iterates $f^n(z)$ do not converge to ∞ for any $z\in U$ [7, Theorem 1], hence U is not a Baker domain. This means that U is either a component of an attracting or parabolic basin, a Siegel disk or a wandering domain [1, Theorem 6]. Assume that U is a wandering domain and let $z \in U$. Then $f^n(z)$ accumulates at points in $\mathcal{J}(f) \cap P(f)$ [2, Theorem] but this set is empty. Hence U is not a wandering domain. For the same reason, U is not a Siegel disk or a component of a parabolic basin (see [1, Theorem 7]), hence U is a component of a basin of attraction of an attracting periodic orbit. Hence $\mathcal{F}(f) = \mathcal{A}(f)$ and every singular value converges under iteration to an attracting periodic orbit.

Now let f be nonescaping-hyperbolic. We have shown that only finitely many attracting cycles contain a singular value in their basins, and since every attracting cycle must intersect $S(f)$ [1, Theorem 7], it follows that $\operatorname{Attr}(f)$ is finite. $\square$

Note that the proof of Proposition 2.2 implies that for a transcendental entire map, hyperbolicity and nonescaping-hyperbolicity are equivalent. Furthermore, a polynomial f is hyperbolic but not nonescaping-hyperbolic if and only if $S(f) \cap I(f) \neq \emptyset$.

By Definition 2.1, every (nonescaping-)hyperbolic transcendental entire function belongs to the *Eremenko-Lyubich class*

$$\mathscr{B} := \{f : \mathbb{C} \to \mathbb{C} \text{ transcendental entire} : S(f) \text{ is bounded}\}.$$

Finally, we would like to remark that an entire map for which every singular value is absorbed by an attracting cycle does not necessarily have dynamics similar to a hyperbolic function. The requirement that $S(f)$ is a bounded set is indeed crucial for Proposition 2.2 to hold: As Example D in [12] shows, there is a transcendental entire map f with wandering domains for which all singular values are mapped by f to an attracting fixed point.

Hausdorff and kernel convergence. Let M be a metric space. Recall that the *Hausdorff distance* between two compact sets $A, B \subset M$ is defined to be

$$d_H(A,B) := \inf\{\varepsilon > 0 : A \subset U_\varepsilon(B),\ B \subset U_\varepsilon(A)\}.$$

For studying convergence of open connected subsets of M we will define a concept analogous to the *Carathéodory* or *kernel convergence* for domains in the complex plane. For more details, see [8, §5].

Definition 2.3 (Kernel)**.** Let $o \in M$ and let $O_n \subset M$, $n \in \mathbb{N}$, be open connected sets containing o. The *kernel* of the sequence (O_n) (w.r.t. o) is the largest open connected set $O \ni o$ such that each compact set $K \subset O$ is contained in all but finitely many O_n.

We call the point o the *marked point* of the sequence (O_n). Clearly, the existence of a kernel is equivalent to the existence of a neighbourhood of o which is contained in all but finitely many O_n.

We say that the sequence (O_n) *converges* to O (*as kernels*) and write $O_n \to O$ if O is a kernel of each subsequence of (O_n). The middle example in Figure 1 shows a sequence that does not converge to its kernel.

Observe that a sequence (O_n) can have more than one kernel (see the left-hand example in Figure 1), and each of them is specified by the choice of a marked point. Now let M



FIGURE 1. *left:* The interiors of the curves are simply-connected domains with three different kernels K_1, K_2 and K_3. *middle:* The domains $D_n := \{z : |z| < 2 - r_n \text{ where } n \equiv r_n \mod 2, r_n \in \{0,1\}\}$ have a unique kernel D_1 but they do not converge to it. *right:* The domains $D_n := \mathbb{C}\setminus(\{z = iy : |y| \geq 1/n\} \cup \{z : |z - n| \leq 1\})$ converge as kernels to the left half-plane $\mathbb{H}_{<0}$ w.r.t. the marked point -1 but their complements do not converge to $\mathbb{C} \setminus \mathbb{H}_{<0}$ in the Hausdorff metric.

be a locally compact metric space, e.g., an analytic manifold, and let O be a kernel of O_n. Since, by definition, every kernel is open and since M is locally compact, every point in O has a compact neighbourhood which is contained in O. This means that we can choose any point $o \in O$ to be the marked point. Hence we will talk about the sets O_n and O without mentioning the marked point if it is implicit from the context which point is meant.

We have the following relation between Hausdorff and kernel convergence.

Proposition 2.4. *Let K_n, K be nonempty compact subsets of a locally compact metric space M. Then $d_H(K_n, K) \to 0$ as $n \to \infty$ if and only if the following two conditions hold:*

- *every component O of $K^c := M\setminus K$ is a kernel of a sequence of components O_n of $K_n^c := M\setminus K_n$,*
- *every kernel of an infinite sequence (O_{n_k}) of components of $K_{n_k}^c$ is a component of K^c.*

We will omit the proof since it is elementary and follows mainly from the definitions of kernel and Hausdorff convergence, and since we do not require the statement for any proof in this article; its purpose is more the illustration of how the given concepts of convergence relate to each other. The right-hand example in Figure 1 shows the necessity of the second requirement in Proposition 2.4.

Dynamical approximation. Let Hol($\mathbb{C}$) be the space of all entire functions. We denote the *locally uniform distance* between $f, g \in \text{Hol}(\mathbb{C})$ by $\chi_{\text{luc}}(f, g)$. The metric $\chi_{\text{luc}}(f, g)$ induces the topology of locally uniform convergence on Hol($\mathbb{C}$), so we say that f_n converge to f *locally uniformly* if and only if $\chi_{\text{luc}}(f_n, f) \to 0$ as $n \to \infty$. (For more details see e.g. [18, Chapter 3].) It follows from the Weierstraß Approximation Theorem [18, Theorem 1.4] that the space Hol($\mathbb{C}$) is closed with respect to this topology.

For entire maps with a non-empty set of singular values we introduce a new metric which combines locally uniform convergence with controlled behaviour on the set of singular values. Hence this metric will be more convenient for the study of dynamics of entire functions. Recall that $\mathrm{Hol}^*(\mathbb{C})$ denotes the set of all entire functions which are not constant or linear.

Definition and Proposition 2.5. *The map* $\chi_{\mathrm{dyn}} : \mathrm{Hol}^*(\mathbb{C}) \to [0, \infty)$ *with*

$$\chi_{\mathrm{dyn}}(f, g) := \chi_{\mathrm{luc}}(f, g) + d_H(S(f), S(g))$$

is a metric, where $d_H(S(f), S(g))$ *is measured with respect to the spherical metric.*

We will say that the sequence f_n *approximates* f dynamically *if* $\chi_{\mathrm{dyn}}(f_n, f) \to 0$ *as* $n \to \infty$.

Maps which are close in the metric χ_{luc} do not necessarily have the property that their sets of singular values are close in the Hausdorff metric. This does not have to be true even in the case of a family of functions which depends holomorphically on some parameter λ, as the following example shows.

Example. Let $f_\lambda(z) = \mathrm{e}^{-\lambda z^2 + z - 2}$ with $\lambda \in \mathbb{C}$. The map $f_0(z) = \mathrm{e}^{z-2}$ has 0 as its only singular value. Now let $\lambda \neq 0$ be any complex number. Then apart from the asymptotic value 0, the map f_λ has the additional critical value $v_\lambda := \mathrm{e}^{1/(4\lambda) - 2}$. Clearly, $v_\lambda \to +\infty$ when $\lambda \searrow 0$.

It is well-known that if f_n is a sequence of entire maps such that $\chi_{\mathrm{luc}}(f_n, f) \to 0$, then for every $w \in S(f)$ there is a sequence $\{w_n : w_n \in S(f_n) \text{ for every } n\}$ such that $w_n \to w$ (see e.g. [11, Theorem 2]), yielding lower semi-continuity for the sets of singular values in case of locally uniform convergence. Hence convergence in the metric χ_{dyn} makes in particular sure that there are no sequences of singular values of the approximating maps f_n that accumulate outside $S(f)$.

Nonescaping-hyperbolicity is not an open property in the topology of locally uniform convergence. For instance, let $f_\lambda(z) := \mathrm{e}^{\lambda z} \cdot z^2$. Then $f_0(z) = z^2$ is nonescaping-hyperbolic while for any sufficiently small $\lambda > 0$, the critical value $4/(\mathrm{e}^2 \lambda^2)$ escapes to ∞. However, the set of nonescaping-hyperbolic entire maps is open in the topology induced by the metric χ_{dyn}.

Theorem 2.6 (Nonescaping-hyperbolicity is an open property)**.** *The set*

$$\mathcal{H} := \{f \in \mathrm{Hol}^*(\mathbb{C}) : f \textit{ is nonescaping-hyperbolic}\}$$

is open in the topology induced by the metric χ_{dyn}.

Note that $\mathcal{H} \subset \mathrm{Hol}^*_b(\mathbb{C})$. Theorem 2.6 will follow from the following lemma.

Lemma 2.7. *Let* $f \in \mathrm{Hol}^*(\mathbb{C})$ *and let* $K \subset \mathcal{A}(f)$ *be a compact set. Then* $K \subset \mathcal{A}(g)$ *for all* $g \in \mathrm{Hol}^*(\mathbb{C})$ *that are sufficiently close to* f *in the metric* χ_{luc}.

Proof. The components of $\mathcal{A}(f)$ form an open cover of the compact set K, hence there is a finite subcover. We can assume w.l.o.g. that K is contained in the basin of attraction

$A(z_0)$ of a fixed point $z_0 \in \mathbb{C}$ of f, since otherwise we can repeat the argument for every attracting periodic point of f. There exists a bounded open set $U \ni z_0$ such that $f(U) \Subset U$. By definition of χ_{luc}, $g(U) \Subset U$ holds for all entire maps g for which $\chi_{\mathrm{luc}}(f, g)$ is sufficiently small. By Montel's Theorem, $\{g^k\}_{k\in\mathbb{N}}$ is normal in U and by the Contraction Mapping Theorem, g has a fixed point in U which is necessarily attracting. Since U is bounded, it follows that $U \subset \mathcal{A}(g)$.

There exists $N \in \mathbb{N}$ such that $f^N(K) \subset U$, so again, by locally uniform convergence, $g^N(K) \subset U \subset \mathcal{A}(g)$ if $\chi_{\mathrm{luc}}(f, g)$ sufficiently small. The claim now follows from the complete invariance of $\mathcal{A}(g)$ under the map g. □

Proof of Theorem 2.6. Let f be nonescaping-hyperbolic, hence $S(f)$ is bounded and contained in $\mathcal{A}(f)$. Choose $\delta > 0$ sufficiently small such that $K := \overline{U_\delta(S(f))} \subset \mathcal{A}(f)$. Since K is compact, it follows from Lemma 2.7 that there is a constant $\varepsilon > 0$ such that $K \subset \mathcal{A}(g)$ for all g with $\chi_{\mathrm{luc}}(f, g) < 2\varepsilon$. Now choose $\eta = \min\{\varepsilon, \delta\}$. Then for all maps g with $\chi_{\mathrm{dyn}}(f, g) < \eta$ we obtain

$$S(g) \subset U_\delta(S(f)) \Subset \mathcal{A}(g),$$

hence g is nonescaping-hyperbolic. □

Remark. We have shown that nonescaping-hyperbolicity is not an open property in the topology of locally uniform convergence. The given example also shows that — in the same topology — the set of all hyperbolic maps in $\mathrm{Hol}^*(\mathbb{C})$ is not open either. Now let p be a hyperbolic polynomial for which a finite singular value escapes to infinity. It is plausible that for any sufficiently small ε, the neighbourhood $U_\varepsilon(p) = \{f \in \mathrm{Hol}^*(\mathbb{C}) : \chi_{\mathrm{dyn}}(f, p) < \varepsilon\}$ of p contains transcendental entire functions with the same property, i.e., maps for which some singular value escapes. Hence it is likely that hyperbolicity is not open in the topology induced by the metric χ_{dyn} either.

3. STABILITY OF NONESCAPING-HYPERBOLIC PARAMETERS

Recall that our goal is to prove that under certain conditions, a kernel of a sequence of nonescaping-hyperbolic components equals a nonescaping-hyperbolic component of the limit family. As we will see, it is not hard to show that every such component of $\mathscr{F}_\infty$ is contained in a kernel of a sequence of nonescaping-hyperbolic components of $\mathscr{F}_n$. This statement requires even less restrictions than those stated in Theorem 1.1. For the other inclusion to hold, we have to construct a more sensible setup.

Denote by $\widehat{\mathbb{N}} := \mathbb{N} \cup \{\infty\}$ the one-point compactification of the set of natural numbers. The metric $\chi_{\widehat{\mathbb{N}}}$ on $\widehat{\mathbb{N}}$ defined by $\chi_{\widehat{\mathbb{N}}}(m, n) := 2^{-\min(m,n)-1}$ when $m \neq n$ (and equal zero otherwise) makes $\widehat{\mathbb{N}}$ a complete metric space. From now on, we assume that M is a complex manifold with a metric χ_M and define $M' := \widehat{\mathbb{N}} \times M$. The relation $\chi_{M'}((m, \lambda), (n, \nu)) := \chi_{\widehat{\mathbb{N}}}(m, n) + \chi_{M'}(\lambda, \nu)$ then defines a metric $\chi_{M'}$ on M'.

For every $n \in \widehat{\mathbb{N}}$ let $\mathscr{F}_n = \{f_{n,\lambda} : \mathbb{C} \to \mathbb{C},\ \lambda \in M\} \subset \mathrm{Hol}^*_b(\mathbb{C})$ be a family of functions parametrized by M. To simplify the notations, we will skip the first index entry for maps in $\mathscr{F}_\infty$, i.e., we will write f_λ for $f_{\infty,\lambda} \in \mathscr{F}_\infty$.

We want all families $\mathscr{F}_n, n \in \widehat{\mathbb{N}}$, to satisfy the following.

Dynamical standing assumption ('dsa')**.** The map

$$F : M' \to \mathrm{Hol}_b^*(\mathbb{C}),\ (n, \lambda) \mapsto f_{n,\lambda}$$

is continuous with respect to the metrics $\chi_{M'}$ and χ_{dyn}.

The key-feature of 'dsa' is the local uniformity in λ *and* n: Let $f_{\lambda_0} \in \mathscr{F}_\infty$ and let $K \subset \mathbb{C}$ be a compact set. Then for every $\varepsilon > 0$ there exist $\delta > 0$ and $n_0 \in \mathbb{N}$ such that $|f_{n,\lambda}(z) - f_{\lambda_0}(z)| < \varepsilon$ for all $z \in K$, $\lambda \in U_\delta(\lambda_0)$ and $n \geq n_0$.

Notations. For every $n \in \widehat{\mathbb{N}}$ we denote by

$$\mathcal{H}(\mathscr{F}_n) := \{\lambda \in M : f_{n,\lambda} \in \mathscr{F}_n \text{ is nonescaping-hyperbolic}\}$$

the parameters corresponding to nonescaping-hyperbolic maps in the respective family. We will usually denote a component of $\mathcal{H}(\mathscr{F}_n)$ by H_n, a component of $\mathcal{H}(\mathscr{F}_\infty)$ by H_∞ and a kernel of a sequence H_n by $\widetilde{H}$.

Proposition 3.1. *Let H_∞ be a component of $\mathcal{H}(\mathscr{F}_\infty)$. Then there exists a kernel $\widetilde{H}$ of a sequence of components of $\mathcal{H}(\mathscr{F}_n)$ such that $H_\infty \subset \widetilde{H}$.*

Proof. Let $\lambda_0 \in \mathcal{H}(\mathscr{F}_\infty)$. It follows from Theorem 2.6 and the dynamical standing assumption that there exists a neighbourhood $U(\lambda_0) \subset M$ such that $U(\lambda_0) \subset \mathcal{H}(\mathscr{F}_n)$ for all sufficiently large $n \in \mathbb{N}$. □

To prove the opposite inclusion, we have to make additional restrictions. Our requirements, that we will assume from now on, are formalized in the following way.

Holomorphic standing assumption ('hsa')**.**

(*i*) For every $n \in \widehat{\mathbb{N}}$, the maps $f_{n,\lambda}$ depend holomorphically on $\lambda \in M$.

(*ii*) Let $n \in \mathbb{N}$ and $\lambda_0 \in M$. Then the singular values of f_{n,λ_0} are holomorphically parametrized by M, i.e., for each singular value s of f_{n,λ_0} there exists a holomorphic map $w_n : M \to \mathbb{C}, \lambda \mapsto w_n(\lambda)$ such that $w_n(\lambda_0) = s$ and $w_n(\lambda) \in S(f_{n,\lambda})$.

Note that the second condition of 'hsa' does not imply that a parametrization of $S(f_{n,\lambda_0})$ for some λ_0 needs to be an exhaustion of the set of singular values for another parameter $\lambda \neq \lambda_0$. A priori, it is possible that $S(f_{n,\lambda_0}) = \{w_n^i(\lambda_0)\}_{i \in I}$ for some index set I but $\{w_n^i(\lambda)\}_{i \in I} \subsetneq S(f_{n,\lambda})$ for some $\lambda \neq \lambda_0$.

Also note that we do not assume condition (*ii*) to be satisfied by the maps in $\mathscr{F}_\infty$. The reason is that we only need a *local* holomorphic parametrization of the sets of singular values for maps in $\mathscr{F}_\infty$ and as the next statement shows, this follows from the assumptions we already made.

Theorem 3.2. *Let $\widetilde{H}$ be a kernel of a sequence of nonescaping-hyperbolic components H_n, $U \ni \lambda_0$ a simply-connected neighbourhood of λ_0 with $U \Subset \widetilde{H}$, and let s be a singular value of f_{λ_0}.*

Then there exists a holomorphic map $w : U \to \mathbb{C}$ such that $w(\lambda) \in S(f_\lambda)$, $w(\lambda_0) = s$ and the family $\{f_\lambda^n(w(\lambda))\}_{n \in \mathbb{N}}$ is normal in U.

Proof. Let $f_{\lambda_0} \in \mathscr{F}_\infty$. By [1, Theorem 5] and [18, Theorem 13.1], the map f_{λ_0} has infinitely many repelling periodic points (in its Julia set). Let us pick two such points and denote them by $p(\lambda_0)$ and $q(\lambda_0)$; let n_1 and n_2 be their periods. If D is a disk at $p(\lambda_0)$ such that $p(\lambda_0)$ is the only periodic point of f_{λ_0} of period $\leq n_1$ in $\overline{D}$, then it follows from 'dsa' and Rouché's theorem that there is a neighbourhood $U(\lambda_0)$ of λ_0 and an integer $n_0 \geq 0$ such that for every $\lambda \in U(\lambda_0)$ and every $n \geq n_0$, the map $f_{n,\lambda}$ has exactly one periodic point $p_n(\lambda)$ of period n_1 in D and no other periodic point of period $\leq n_1$ in $\overline{D}$. By the Cauchy Integral Formula (after decreasing the initial disk D, if necessary), every such periodic point $p_n(\lambda)$ must be repelling. By the Implicit Function Theorem, every of these points can be analytically continued as a repelling periodic point of period n_1 in a sufficiently small neighbourhood. The previous observation then implies that for every sufficiently large $n \in \widehat{\mathbb{N}}$, there exists an analytic function $p_n : U(\lambda_0) \to \mathbb{C}$ such that $p_n(\lambda) \in D$ is a repelling periodic point of $f_{n,\lambda}$ of period n_1. By construction, $p_n(\lambda) \to p(\lambda)$ when $n \to \infty$. We can repeat the same procedure for $q(\lambda_0)$ and obtain holomorphic maps $q_n : U'(\lambda_0) \to \mathbb{C}$.

Let us now assume that $\lambda_0 \in \widetilde{H}$. Since $\widetilde{H}$ is open, there is a disk $B \subset \widetilde{H}$ centred at λ_0 such that the maps p_n, q_n are defined and holomorphic in B and their images are repelling periodic points of the corresponding maps (and periods). Let $U \supset B$ be a simply-connected bounded domain with closure in $\widetilde{H}$. Since $\widetilde{H}$ is a kernel of a sequence H_n of nonescaping-hyperbolic components, the compact set $\overline{U}$ is contained in all H_n for $n \in \mathbb{N}$ chosen sufficiently large. Hence, the maps p_n and q_n ($n \in \mathbb{N}$) can be holomorphically continued to all of U, since otherwise the Implicit Function Theorem would imply that for some $\tilde{\lambda} \in U$, the map $f_{n,\tilde{\lambda}}$ has an indifferent periodic point.

Let us now consider the maps $\Phi_\lambda(z) = \frac{z-p(\lambda)}{q(\lambda)-p(\lambda)}$ and $\Phi_{n,\lambda}(z) = \frac{z-p_n(\lambda)}{q_n(\lambda)-p_n(\lambda)}$. Conjugating f_λ with Φ_λ and $f_{n,\lambda}$ with $\Phi_{n,\lambda}$, we obtain conformal conjugates such that the points 0 and 1 correspond to our previously considered repelling periodic points. Hence we can assume that $p(\lambda) = 0$, $q(\lambda) = 1$ for all $\lambda \in B$ and $p_n(\lambda) = 0$, $q_n(\lambda) = 1$ for all $\lambda \in U$. In particular,

$$\{0,1\} \subset \mathcal{J}(f_\lambda), \mathcal{J}(f_{n,\lambda}) \text{ and } S(f_{n,\lambda}) \subset \mathbb{C}\backslash\{0,1\}.$$

for all sufficiently large integers $n \in \widehat{\mathbb{N}}$ and the corresponding values of λ.

Let s be a singular value of f_{λ_0}. By 'dsa', there is a sequence of singular values s_n of the maps f_{n,λ_0} that converges to s. Due to 'hsa', there are holomorphic maps w_n such that $w_n(\lambda_0) = s_n$ and $w_n(\lambda) \in S(f_{n,\lambda})$ for all $\lambda \in M$. By the previous argument, we have that $w_n(U) \subset \mathbb{C}\backslash\{0,1\}$, hence, by Montel's theorem, $\{w_n\}_{n\in\mathbb{N}}$ is a normal family on U. Let (w_{n_k}) be a convergent subsequence of (w_n), and let w be a limit function which is necessarily holomorphic. By construction we have that $w(\lambda_0) = s$, and 'dsa' implies that $w(\lambda) \in S(f_\lambda)$ holds for all $\lambda \in U$. Hence w is a holomorphic parametrization of the singular value s on U.

Consider now for a fixed ν the family $\{f^\nu_{n,\lambda}(w_n(\lambda))\}_{n\in\mathbb{N}}$ with $\lambda \in U$. Since the Fatou set of an entire map is completely invariant, $w_n(\lambda) \in \mathcal{F}(f_{n,\lambda})$ implies that $f^\nu_{n,\lambda}(w_n(\lambda)) \subset \mathcal{F}(f_{n,\lambda}) \subset \mathbb{C}\backslash\{0,1\}$. Applying Montel's theorem it follows that for each ν the above family is normal in U. For simplicity, denote its limit by $S_\nu : U \to \widehat{\mathbb{C}}$, $\lambda \mapsto S_\nu(\lambda) :=$

$\lim_{n\to\infty} f^{\nu}_{n,\lambda}(w_n(\lambda))$. It follows from the local uniform convergence of the maps f_n and w_n that

$$S_\nu(\lambda) = \lim_{n\to\infty}\lim_{m\to\infty} f^{\nu}_{n,\lambda}(w_m(\lambda)) = \lim_{n\to\infty} f^{\nu}_{n,\lambda}(w(\lambda)) = f^{\nu}_{\lambda}(w(\lambda)).$$

By Hurwitz's theorem, either $S_\nu \subset \mathbb{C}\backslash\{0,1\}$ or $S_\nu \equiv 0$ or 1. Recall that 0 and 1 are periodic points of f_λ, hence if $S_\nu \equiv 0$ (resp. 1) then there exists some $m \in \mathbb{N}$ such that $S_{\nu+km} \equiv 0$ (resp. 1) for all $k \in \mathbb{N}$. Applying Montel's theorem once more, we obtain that $\{S_\nu\}_{\nu\in\mathbb{N}}$ is a normal family on U. □

We can now prove the remaining statement which will then imply Theorem 1.1.

Theorem 3.3. *Let $\widetilde{H}$ be a kernel of a sequence of components of $\mathcal{H}(\mathscr{F}_n)$. If there is a component H_∞ of $\mathcal{H}(\mathscr{F}_\infty)$ such that $\widetilde{H} \cap H_\infty \neq \emptyset$, then $\widetilde{H} \subset H_\infty$.*

Proof. We will prove the statement by contradiction, so assume that there exists some $\lambda_0 \in \widetilde{H} \cap \partial H_\infty$. With the same notations as in the previous proof, it follows that $S_\nu(B) \subset \mathbb{C}\backslash\{0,1\}$ for all $\nu \in \mathbb{N}$, since $B' := B \cap H_\infty \neq \emptyset$ (and hence $S_\nu|_B$ cannot be constant 0 or 1).

For a limit function S of S_ν we have that either $S \equiv c \in \widehat{\mathbb{C}}$, in which case $S \equiv \infty$, or S is a non-constant function with $S(B) \subset \mathbb{C}\backslash\{0,1\}$.

The case $S \equiv \infty$ clearly cannot occur since this would mean that some singular value converges to ∞ for all parameters in B but the nonempty subset B' of B consists of nonescaping-hyperbolic parameters.

Let $S_\nu(B) \subset \mathbb{C}\backslash\{0,1\}$. For all $\lambda \in B'$ there is a holomorphic map $a : B' \to \mathbb{C}$ such that $a(\lambda)$ is an attracting point of f_λ which attracts $w(\lambda)$. Since $B \supset B'$ is simply-connected, the point $a(\lambda)$ can then be analytically continued to an attracting point on the entire domain B. The image $a(B)$ is bounded, hence for every $\lambda \in B$ the singular value $w(\lambda)$ is attracted by a finite attracting periodic cycle of f_λ. Since $s = w(\lambda_0)$ was assumed to be an arbitrary singular value of f_{λ_0}, we can repeat this procedure for any of the singular values of f_{λ_0}. Recall that, by assumption, $\mathscr{F}_\infty \subset \mathrm{Hol}^*_b(\mathbb{C})$, hence the singular sets are bounded. This implies that $B \ni \lambda_0$ is contained in some nonescaping-hyperbolic component H_∞ of $\mathscr{F}_\infty$, contradicting the assumption that $\lambda_0 \in \widetilde{H} \cap \partial H_\infty$. □

Without the assumption of dynamical approximation, (which is part of 'dsa') we cannot expect that a kernel $\widetilde{H}$ is always a nonescaping-hyperbolic component of the family $\mathscr{F}_\infty$. It is easy to find suitable examples. One such example was given in [14]: the authors approximated a holomorphic family of quadratic polynomials by families of polynomials of degree four, such that a kernel of a sequence of nonescaping-hyperbolic components was a proper subset of some nonescaping-hyperbolic component of the limit family.

Here we give an example which respects our standing assumptions, showing that the case $\widetilde{H} \cap \mathcal{H}(\mathscr{F}_\infty) = \emptyset$ in Theorem 1.1 does indeed occur.

Example. Let

$$P_{\lambda,n}(z) = z^3 - \left(\frac{\lambda - 1}{\sqrt{\mu_n + \lambda - 2}} + \sqrt{\mu_n + \lambda - 2}\right) z^2 + \lambda z, \tag{1}$$

where $\lambda \in \mathbb{C} \setminus [1, 5]$, $|\mu_n| < 1$ does not depend on λ and $\mu_n \to 1$ as $n \to \infty$ (for instance we can choose $\mu_n = \frac{n}{n+1}$). Note that $P_{\lambda,n}(z)$ and $P_\lambda(z) = z^3 - 2\sqrt{\lambda - 1}z^2 + \lambda z$ satisfy 'dsa' and 'hsa'. Every $P_{\lambda,n}$ has 0 as a fixed point of multiplier λ. Furthermore, there is a second fixed point $a_n = \sqrt{\mu_n + \lambda - 2}$ with multiplier μ_n. Thus, if we choose $\lambda \in \mathbb{D}$ then every polynomial $P_{\lambda,n}$ is nonescaping-hyperbolic. Hence there is a kernel $\widetilde{H}$ of components of $\mathcal{H}(\mathscr{F}_n)$ which contains the unit disk $\mathbb{D}$. On the other hand, every P_λ has a parabolic fixed point at $a = \sqrt{\lambda - 1}$, hence $\mathcal{H}(\mathscr{F}_\infty) = \emptyset$.

Nevertheless, the behaviour of f_λ is still stable in the sense of *J-stability* for parameters belonging to a kernel. Here, J-stability is defined analogously to the case of rational maps or transcendental entire maps with finitely many singular values (see e.g. [7]):

Definition 3.4. Let $\mathscr{F} = \{f_\lambda : \mathbb{C} \to \mathbb{C} : \lambda \in M\}$ be a holomorphic family of entire functions. A map $f_{\lambda_0} \in G$ is said to be *J-stable* if $\mathcal{J}(f_{\lambda_0})$ moves holomorphically in a neighbourhood $\Lambda \subset M$ of λ_0, i.e., if there is a holomorphic motion

$$\Phi : \Lambda \times \mathcal{J}(f_{\lambda_0}) \to \mathbb{C}$$

such that

$$\Phi(\lambda, \mathcal{J}(f_{\lambda_0})) = \mathcal{J}(f_\lambda) \text{ and } \Phi(\lambda, f_{\lambda_0}(z)) = f_\lambda(\Phi(\lambda, z))$$

for all $z \in \mathcal{J}(f_{\lambda_0})$ and all $\lambda \in \Lambda$.

Φ being a holomorphic motion means that Φ is injective in z when λ is fixed, holomorphic in λ for fixed z and that $\Phi_{\lambda_0} \equiv \mathrm{id}$. For more details see [7, Section 8].

The map Φ_λ is a conjugacy between f_{λ_0} and f_λ on their Julia sets, hence it maps periodic points of f_{λ_0} to periodic points of f_λ. Since periodic points form a dense subset of the Julia set, such a conjugacy is unique if it exists.

Theorem 3.5. *Let $\widetilde{H}$ be a kernel of a sequence of components of $\mathcal{H}(\mathscr{F}_n)$ and let $\lambda \in \widetilde{H}$. Then $f_\lambda \in \mathscr{F}_\infty$ is J-stable.*

Proof. We will prove the contraposition, so let λ_0 be a parameter for which f_{λ_0} is not J-stable and let Λ be a simply-connected bounded neighbourhood of λ_0. Then there is some repelling periodic point, say of period n, of f_{λ_0} which has no analytic continuation as a solution of the equation $f_\lambda^n(z) - z = 0$. Otherwise, it would follow from the λ-lemma [7] that the closure of all repelling points of f_{λ_0}, which equals the Julia set of f_{λ_0}, would move holomorphically, contradicting that f_{λ_0} is not J-stable.

Let $p(\lambda_0)$ be such a repelling periodic point and let $\gamma : [0, 1] \to \Lambda$, $\gamma(0) = \lambda_0$, be a path along which $p(\lambda_0)$ cannot be continued analytically. By the Implicit Mapping Theorem, the point $p(\lambda)$ for $\lambda = \gamma(1)$ must be indifferent. By the Minimum Modulus Principle we can then find a nearby path $\tilde{\gamma} \subset \Lambda$ connecting λ_0 to some parameter $\tilde{\lambda}$ along which the considered point becomes attracting. Hence there is a singular value $s(\tilde{\lambda})$ of $f_{\tilde{\lambda}}$ converging to the attracting periodic point $p(\tilde{\lambda})$.

Now let us assume that our assumption is wrong, meaning that λ_0 belongs to some kernel $\widetilde{H}$. We can assume w.l.o.g. that the neighbourhood Λ was chosen sufficiently small such that $\Lambda \Subset \widetilde{H}$. Then Theorem 3.2 implies that there is a holomorphic parametrization w of the singular value $s(\tilde{\lambda})$ such that $\{f_\lambda^n(w(\lambda))\}$ is a normal family on Λ. But then each $w(\lambda)$ converges to an attracting point $p(\lambda)$ of f_λ, in which case $p(\lambda)$ can be continued analytically to an attracting point of f_λ on the whole of Λ, contradicting the fact that $p(\lambda_0)$ is repelling. ☐

4. Construction of examples

As we have seen in the previous section, a sequence of families to which our results apply has to satisfy two primary conditions which we have formulated as the standing assumptions 'dsa' and 'hsa'. Hence starting with an entire map $f \in \mathrm{Hol}_b^*(\mathbb{C})$, we can split the problem into the following two:

(*i*) Find a sequence f_n of entire functions which approximates f *dynamically.*
(*ii*) Construct *holomorphic* families $\{f_{n,\lambda}\}$ and $\{f_\lambda\}$ using the functions f_n, f.

We will start with the second problem. It turns out that there is a natural way to find suitable holomorphic families for *any* entire function.

Holomorphic families inside quasiconformal equivalence classes. Recall that M is a complex manifold. Let μ be a k-Beltrami coefficient of $\mathbb{C}$, i.e., $\mu : \mathbb{C} \to \mathbb{C}$ is a measurable function such that $\|\mu\|_\infty \leq k < 1$ almost everywhere (a.e.) in $\mathbb{C}$. By the Integrability Theorem [15, Theorem 4.4], there exists a k-quasiconformal homeomorphism $\Psi : \mathbb{C} \to \mathbb{C}$ whose complex dilatation equals μ a.e. in $\mathbb{C}$.

Let $f \in \mathrm{Hol}_b^*(\mathbb{C})$ be an entire map. Then the *pull-back* $f^*\mu$ of μ by f is given by

$$f^*\mu(z) := (\mu \circ f)\frac{\overline{f'(z)}}{f'(z)}.$$

In particular, $\|\mu\|_\infty \leq k$ implies that $\|f^*\mu\|_\infty \leq k$.

By the Integrability Theorem there exists a quasiconformal homeomorphism Φ whose complex dilatation equals $f^*\mu$ a.e. in $\mathbb{C}$. A formal computation yields that the complex dilatation of the map $g = \Psi \circ f \circ \Phi^{-1}$ is 0 a.e., which means that g is a holomorphic map [15, Theorem 1.1]. We say that g is *quasiconformally equivalent* to f.

Let $\Lambda \subset M$ be an open connected set and let $\{\Psi_\lambda\}$ be a family of quasiconformal homeomorphisms with uniformly bounded complex dilatations that depend holomorphically on $\lambda \in \Lambda$. The parametrized version of the Integrability Theorem [10, Chapter 4.7] gives the following way to construct such a family, starting from a Beltrami coefficient μ_0:

Let $(\mu_\lambda)_{\lambda\in\Lambda}$ be a holomorphic family of Beltrami coefficients with $\|\mu_\lambda\|_\infty \leq k < 1$ that contains μ_0, i.e., $\mu_0 \equiv \mu_{\lambda_0}$ for some $\lambda_0 \in \Lambda$. (One way to embed μ_0 in a holomorphic family of Beltrami coefficients is as follows: Let $h : \Lambda \to D_r(0)$ be a holomorphic map, where r is chosen such that $(1 + r) \cdot \|\mu_0\|_\infty < 1$ and $h(\lambda_0) = 0$. Then the functions

$\mu_\lambda := (1+h(\lambda))\cdot\mu_0$ form a holomorphic family of Beltrami coefficients.) For every λ, let Ψ_λ be a quasiconformal homeomorphism with Beltrami coefficient μ_λ, chosen such that all Ψ_λ have the same parametrization (e.g., all Ψ_λ fix 0, 1 and ∞). Then the Integrability Theorem implies that $\lambda \mapsto \Psi_\lambda$ is also holomorphic.

As in the previous construction we obtain a family $\mathscr{F}_\infty = \{f_\lambda = \Psi_\lambda \circ f \circ \Phi_\lambda^{-1}\}_{\lambda\in\Lambda}$ of entire maps. By computing the derivative of the equation $\Psi_\lambda \circ f = f_\lambda \circ \Phi_\lambda$ with respect to $\overline{\lambda}$, we get that the functions f_λ also depend holomorphically on λ (see e.g. the proof of Proposition 13 in [3]). Since an entire function f is a covering map when restricted to $\mathbb{C}\backslash f^{-1}(S(f))$, we obtain that $S(f_\lambda) = \Psi_\lambda(S(f))$, hence the set of singular values moves holomorphically in the family $\mathscr{F}_\infty$ in a unique way. Furthermore, $S(f_\lambda)$ is bounded for all λ since the maximal dilatations of the quasiconformal maps Ψ_λ are uniformly bounded.

Let (f_n) be a sequence of entire functions which converges to f dynamically. In the same manner as above, we define for each n the holomorphic family $\mathscr{F}_n = \{f_{n,\lambda} = \Psi_\lambda \circ f_n \circ \Phi_{n,\lambda}^{-1}\}_{\lambda\in\Lambda}$ by starting with the same family $\{\Psi_\lambda\}$ of quasiconformal homeomorphisms. As before, the set of singular values $S(f_{n,\lambda}) = \Psi_\lambda(S(f_n))$ moves holomorphically. Hence it remains to show that the map

$$F : M' \to \mathrm{Hol}_b^*(\mathbb{C}),\ (n,\lambda) \mapsto f_{n,\lambda}$$

is continuous. In other words, let $\lambda_0 \in M$ and let $(n,\lambda) \to (\infty,\lambda_0)$. We have to show that in this case,

$$\chi_{\mathrm{luc}}(f_{n,\lambda}, f_{\lambda_0}) + d_H(S(f_{n,\lambda}), S(f_{\lambda_0})) \to 0.$$

Clearly, $d_H(S(f_{n,\lambda}), S(f_{\lambda_0})) = d_H(\Psi_\lambda(S(f_n)), \Psi_{\lambda_0}(S(f))) \to 0$ as $(n,\lambda)\to(\infty,\lambda_0)$, since Ψ_λ depends holomorphically (so in particular continuously) on λ and the maps f_n approximate f dynamically.

To see that $\chi_{\mathrm{luc}}(f_{n,\lambda}, f_{\lambda_0}) \to 0$, we have to look at the sequence of pull-backs

$$f_n^*\mu_\lambda = (\mu_\lambda \circ f_n)\frac{\overline{f_n'(z)}}{f_n'(z)},$$

where μ_λ denotes the complex dilatation of Ψ_λ. By assumption, there exists a constant $k<1$ such that $\|f_n^*\mu_\lambda\|_\infty < k < 1$ for all n. Since also $f_n^*\mu_\lambda \to f^*\mu_{\lambda_0}$ a.e., it follows that the (uniquely normalized solutions) $\Phi_{n,\lambda}$ converge locally uniformly to Φ_{λ_0} [15, Theorem 4.6], yielding the desired statement.

Functions of sine type. The above construction yields many examples of families where Theorems 3.2 and 3.3 can be applied to, provided that we have a sequence of functions approximating f dynamically. There are clearly various ways of approximating an entire function locally uniformly but it is a very strong requirement to keep control over the sets of singular values, since this set can be arbitrarily complicated (for instance, it can have nonempty interior).

For certain (families of) transcendental entire functions that were and are of particular interest in holomorphic dynamics, appropriate approximations are known and were extensively studied. For instance, one can approximate the exponential map by the polynomials

$P_n(z) = (1 + z/n)^n$ or the function $f(z) = \sin(z)/z$, which has infinitely many singular values, by the sequence $T_n(z)/z$, where $T_n(z)$ denotes the Chebyshev polynomial of degree n.

We want to introduce another set of transcendental entire functions for which a dynamical approximation exists, namely *real sine-type maps with real zeros.*

Definition 4.1. A function f is said to be of *sine type* σ if there are positive constants c, C, τ such that

$$c\,\mathrm{e}^{\sigma|\operatorname{Im} z|} \leq |f(z)| \leq C\,\mathrm{e}^{\sigma|\operatorname{Im} z|},$$

where the upper estimate holds everywhere in $\mathbb{C}$ and the lower estimate holds at least outside the horizontal strip $|\operatorname{Im} z| \leq \tau$.

Let f be a sine-type function and denote by z_n the zeros of f. By [16, Lecture 17] or ([20, Theorem 3]), the limit

$$\lim_{R\to\infty} \prod_{|z_n|\leq R} \left(1 - \frac{z}{z_n}\right)$$

exists uniformly on compact subsets of $\mathbb{C}$, and it defines an entire function called the *generating function of the sequence* (z_n) which equals f up to $K \cdot z^m$ where K is a constant and $m \geq 0$ is an integer [20, Theorem 2]. By definition, the zeros of f are contained in a horizontal strip around the real axis and f has exactly two tracts over ∞, each of which contains some upper and lower halfplane, respectively. It follows from the Ahlfors-Denjoy Theorem [19, XI, §4, 269] that f has at most two asymptotic values. The derivative of the generating function $\tilde{f}$ of (z_n) is given by $\tilde{f}' = \tilde{f} \cdot \sum \frac{1}{z-z_n}$ and an elementary computation shows that $\tilde{f}$ and hence f has no critical points outside a sufficiently wide horizontal strip. So the set of critical values of f is bounded, implying that every f belongs to the Eremenko-Lyubich class $\mathscr{B}$.

It is now easy to show the following.

Proposition 4.2. *Let f be a real sine-type function for which all zeros are real. Then there exists a sequence p_n of polynomials such that $\chi_{\mathrm{dyn}}(p_n, f) \to 0$ when $n \to \infty$.*

Proof. By a theorem of Laguerre [21, Chapter 8.51], all zeros of $f'(z)$ are real and are separated from each other by the zeros of f. By basic calculus arguments, the generating polynomials cannot have "free" critical values, and by [11, Theorem 2] each singular value of f is approximated by a sequence of singular values of the generating polynomials. Hence the sets of singular values converge in the Hausdorff metric. □

Remark. Standard approximation methods in function theory do not relate to the sets of singular values in a way that would enable us to construct a sequence of entire maps that dynamically approximate a given function f in a nontrivial way. This leaves open an interesting function-theoretic question.

References

[1] W. Bergweiler, 'Iteration of meromorphic functions', *Bull. Amer. Math. Soc.* 29, no. 2 (1993), 151–188, arXiv:math.DS/9310226.

[2] W. Bergweiler, M. Haruta, H. Kriete, H. G. Meier and N. Terglane, 'On the limit functions of iterates in wandering domains', *Ann. Acad. Sci. Fenn., Ser A I Math.* 18 (1993), 369–375.

[3] X. Buff and A. Cheritat, 'Upper bound for the size of quadratic Siegel disks', *Inventiones Mathematicae* 156, no. 1 (2004), 1–24.

[4] R. L. Devaney, L. R. Goldberg and J. H. Hubbard, 'A dynamical approximation to the exponential map by polynomials', *Preprint, Berkley* (1986).

[5] C. Bodelón, R. L. Devaney, M. Hayes, G. Roberts, L. R. Goldberg and J. H. Hubbard, 'Dynamical convergence of polynomials to the exponential', *Journal of Difference Equations and Applications* 6 (2000), 275–307.

[6] A. E. Eremenko, 'On the iteration of entire functions', *Dynamical Systems and Ergodic Theory, Proc. Banach Center Publ. Warsaw* 23 (1989), 339–345.

[7] A. E. Eremenko and M. Y. Lyubich, 'Dynamical properties of some classes of entire functions', *Ann. Inst. Fourier, Grenoble* 42, no. 4 (1992), 989–1020.

[8] G. M. Golusin, 'Geometrische Funktionentheorie', *VEB Deutscher Verlag Der Wissenschaften* (1957).

[9] L. R. Goldberg and L. Keen, 'A finiteness theorem for a dynamical class of entire functions', *Ergodic Theory Dynamical Systems* 6 (1986), 183–192.

[10] J. H. Hubbard, 'Teichmüller theory and applications to geometry, topology and dynamics Volume I: Teichmüller theory. Vol. 1', *Matrix Editions, Ithaca, New York* (2006).

[11] M. Kisaka, 'Local uniform convergence and convergence of Julia sets', *Nonlinearity* 8 (1995), 273–281.

[12] M. Kisaka and M. Shishikura, 'On multiply connected wandering domains of entire functions', in: Transcendental dynamics and complex analysis (P. J. Rippon and G. M. Stallard, eds.), no. 348, *London Mathematical Society Lecture Notes Series, Cambridge University Press* (2008).

[13] B. Krauskopf and H. Kriete, 'Kernel convergence of hyperbolic components', *Ergodic Theory Dynamical Systems* 17 (1997), 1137–1146.

[14] B. Krauskopf and H. Kriete, 'On the convergence of hyperbolic components in families of finite type', *Mathematica Gottingensis - Schriftenreihe des Mathematischen Instituts der Universität Göttingen* (1995).

[15] O. Lehto, 'Univalent Functions and Teichmüller Spaces', *Graduate Texts in Mathematics 109, Springer-Verlag New York, Berlin, Heidelberg* (1986).

[16] B. Ya. Levin, 'Lectures on entire functions', *Translations of Mathematical Monographs, American Mathematical Society* 150, no. 150 (1996).

[17] R. Mañé and P. Sad and D. Sullivan, 'On the dynamics of rational maps', *Annales scientifiques de l'École Normale Supérieure (Sér. 4)* 16, no. 2 (1983), 193–217.

[18] J. Milnor, 'Dynamics in one complex variable', *Annals of Mathematics Studies, Princeton University Press, Princeton, NJ, Vol. 160, 3rd edition* (2006).

[19] R. Nevanlinna, 'Eindeutige analytische Funktionen', *Springer-Verlag/Berlin, Göttingen, Heidelberg, 2nd edition* (1953).

[20] G. Semmler, 'Complete interpolating sequences, the discrete Muckenhoupt condition, and conformal mapping', *Preprint* (2007), arXiv:math.DS/0708.2787.

[21] E. C. Titchmarsh, 'The Theory of Functions', *Oxford University Press* (1932).